

# **Séminaire UMR Sysiphe :**

**Lundi 13 septembre, de 13h à 14 h :**

**Lieu : Université Paris VI, 4 place Jussieu, Paris 5°, métro Jussieu  
Tour 56, 3° étage, couloir 56-46, salle "Darcy"**

## **MODELING FLOW AND TRANSPORT THROUGH HIGHLY HETEROGENEOUS FORMATIONS: CONCEPTUAL ASPECTS**

**by Gedeon Dagan**

Faculty of Engineering, Tel Aviv University, Israel

The widely used analytical models of flow and transport through aquifers of spatially variable conductivity are based on a first-order approximation in  $\sigma^2_Y$ , the variance of the log-conductivity  $Y = \ln K$ . With  $Y$  modelled as random, a few flow (effective conductivity, head and velocity covariances) and transport (macrodispersivity) statistical moments depend on the mean and two-point covariance of  $Y$  only. Thus, the structure is essentially characterized by three parameters:  $K_G$  (the geometric mean),  $\sigma^2_Y$  and  $I_Y$  (integral scale, with two different values for horizontal and vertical directions for anisotropic aquifers).

Numerical modelling requires the knowledge of the entire statistical structure of  $Y$  and the traditional approach is to regard it as multi-Gaussian, i.e. the vector of  $Y$  values at different points is a multivariate normal vector. Comparison with first-order results showed that the latter are quite robust and apply to  $\sigma^2_Y$  as large as unity.

Recently, there is considerable interest in the behaviour of highly heterogeneous formations, of  $\sigma^2_Y$  values larger than unity. A few numerical studies were conducted primarily for 2D (two-dimensional) configurations. The multi-Gaussian model was questioned and alternative ones were considered. A brief review is presented, with emphasis on the multi-indicator model developed recently by the author in collaboration with A. Fiori and I. Janković.

The main aim of the presentation is to discuss a few conceptual problems arising in modelling of flow and transport in highly heterogeneous aquifers such as: the non-uniqueness of structure models in view of the scarcity of field data and criteria of selection of models; 2D versus 3D models; effective conductivity, stationarity and ergodicity requirements; macrodispersivity, its meaningfulness; robustness of first-order approximation and conditioning on data.